

Available online at www.sciencedirect.com**ScienceDirect****Biomedical Journal**journal homepage: www.elsevier.com/locate/bj**Review Article: Special Edition****Purinergic signaling and infection by *Leishmania*:
A new approach to evasion of the immune
response****Amanda Braga de Figueiredo^a, Miriam Conceicao Souza-Testasica^b,
Luis Carlos Crocco Afonso^{a,*}**^a Immunoparasitology Laboratory, Department of Biological Sciences, ICEB/NUPEB, Federal University of Ouro Preto, Brazil^b Coordination Area of Biological Sciences, Federal Institute of Minas Gerais, Brazil

CrossMark

Prof. Luis Carlos
Crocco Afonso**ARTICLE INFO****Article history:**

Received 24 March 2016

Accepted 2 August 2016

Available online 21 September 2016

Keywords:*Leishmania*

Innate response

Immune evasion

Purinergic signaling

ABSTRACT

Infection by protozoan parasites is part of the most common Tropical Neglected Diseases. In the case of leishmaniasis, several millions of people are at risk of contracting the disease. In spite of innumerable studies that elucidated the immune response capable of killing the parasite, the understanding of the evasion mechanisms utilized by the parasite to survive within the very cell responsible for its destruction is still incomplete. In this review, we offer a new approach to the control of the immune response against the parasite. The ability of the parasite to modulate the levels of extracellular ATP and adenosine either by directly acting on the levels of these molecules or by inducing the expression of CD39 and CD73 on the infected cell may influence the magnitude of the immune response against the parasite contributing to its growth and survival.

Leishmaniasis comprises a group of parasitic diseases caused by several species of the *Leishmania* genus. Depending on the combination of the parasite species and yet unknown factors of the host, the disease can manifest itself as a single cutaneous ulcer, multiple lesions throughout the body or even visceral infection which, when untreated, is often lethal [1]. The disease is transmitted by infected females of sand flies of the Phlebotominae family during blood feeding [2] and affects

approximately 300,000 people worldwide annually, mainly in the tropical and subtropical areas [3].

The immune response against the parasite is dependent both on the innate as well as the adaptative responses (for recent reviews, see [4–7]). In this review we will focus primarily on the innate aspects of the immune response and the role of the purinergic signaling in the modulation of such response.

* Corresponding author. Immunoparasitology Laboratory, Department of Biological Sciences, ICEB/NUPEB, Federal University of Ouro Preto, Cruise hill, 35400-000 Ouro Preto, MG, Brazil. Tel.: +55 31 35591701; fax: +55 31 35591680.

E-mail address: afonso@nupeb.ufop.br (L.C.C. Afonso).

Peer review under responsibility of Chang Gung University.

<http://dx.doi.org/10.1016/j.bj.2016.08.004>

2319-4170/© 2016 Chang Gung University. Publishing services by Elsevier B.V. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).

Innate response to *Leishmania*

During blood feeding of infected phlebotomine females, neutrophils rapidly migrate to the site of infection and phagocytize metacyclic promastigotes that are injected together with the insect saliva [8,9]. The role of neutrophils during infection is contradictory. These cells have been reported to help eliminate the parasite via the production of ROS and the induction of extracellular traps [10,11]. However, they also may play a role in inhibiting macrophage activation, due to their death by apoptosis after infection by the parasite [12]. Neutrophils also secrete chemokines that recruit monocytes/macrophages as well as dendritic cells to the site of infection [4,13].

Dendritic cells (DC) are antigen presenting cells capable of producing IL-12 upon infection with some *Leishmania* species such as *Leishmania major* [14]. This production of IL-12 is crucial to the development of a specific Th1 response that is protective to the host [15,16]. However, *Leishmania amazonensis*, which causes a severe form of cutaneous leishmaniasis (diffuse cutaneous leishmaniasis) characterized by the lack of specific immune response in patients [17], inhibits DC activation and interfere with antigen presentation and development of T cell response [18] thus evading the immune response both in humans and in the mouse model [17–21].

Even though *Leishmania* can infect neutrophils and DC, macrophages are the cells in which these parasites live and multiply within the infected mammalian host. *Leishmania* gain access to macrophages via phagocytosis mediated by complement fragments that are deposited and inactivated on the surface of the parasite, due to the action of proteases present on the surface of the infective promastigote [22–25]. The entry of the parasite via complement receptor (CR3)-mediated phagocytosis interferes with macrophage activation allowing the parasite to prevent the initial respiratory burst that, otherwise, would destroy the parasite [26]. However, upon activation by IFN- γ [27,28], produced by both NK [29] and T cells (both CD4 [30] and CD8 [31]), macrophages are activated to produce ROS and NO which are the main effector mechanisms for parasite killing within these cells [32].

In spite of the potential ability of the host to control parasite growth, several mechanisms contribute to evasion of the immune response by *Leishmania*. These mechanisms include, in addition to those already mentioned above, escape from complement activation [26,33], inhibition of macrophage activation by apoptotic neutrophil [12], production of inhibitory cytokines such as IL-10 and TGF- β by macrophages and DC, and the induction of regulatory T cells capable of controlling the immune response [4,19,21]. Although these inhibitory mechanisms have been extensively studied, the evasion of the immune response by some parasite species, in special *L. amazonensis*, is not restricted to these mechanisms. For example, IL-10 deficient mice infected with *L. amazonensis* still develop lesions in spite of an increased Th1 response [34]. Thus, it is conceivable that other mechanisms of immune evasion are present during *Leishmania* infection and modulation of the immune response via purinergic signaling may be one of such mechanisms.

Purinergic signaling and inflammation

During infection or cell injury, ATP can be released to the extracellular milieu and has been described as a danger signal, alerting the immune system to alterations in cellular integrity [35–38]. ATP can also be released by intact cells through connexin and pannexin channels and also via the P2X₇ receptor [39].

Extracellular ATP is a potent inducer of inflammation characterized by macrophage and DC activation and increased production of IL-12 and TNF- α by these cells [40–45]. ATP exerts its effects by binding to P2 receptors which are divided in two subtypes: P2X receptors, that are associated to ionic channels and protein G coupled-P2Y [46–48]. In immune cells, P2X₇ is the main ATP receptor and its activation accounts for most of the inflammatory effects of extracellular ATP [49–51].

In order to regulate the extracellular effects of ATP, its concentration is controlled by the action of extracellular enzymes of which the main players in immune cells are the ecto-NTPDase (CD39) that hydrolyses ATP to ADP and subsequently to AMP and the ecto-5'-nucleotidase (CD73) that removes the phosphate group of AMP leading to the production of adenosine. Adenosine is, then, deaminated to inosine by adenosine deaminase [52,53]. In addition, adenosine concentrations inside and outside the cell are kept relatively constant (between 30 and 300 nM) by the action of bidirectional nucleoside transporters. However, in pathophysiological conditions such as hypoxia, ischemia and cell injury, adenosine extracellular concentrations can peak at 10 μ M [54–56].

By acting on P1 receptors, in particular A_{2A} and A_{2B}, adenosine counteracts the inflammatory effects of ATP [57]. Thus adenosine inhibits the production of inflammatory cytokines by macrophages and DC and the production of microbicidal effectors by neutrophils and macrophages. In addition, adenosine stimulates the synthesis of IL-10, one of the major regulatory cytokines [47,58–63].

Thus, the balance between extracellular ATP and adenosine can contribute to the control of inflammation by, respectively, stimulating or inhibiting the cells involved in the immune response. Next, we present data from the recent literature that show that the purinergic system may interfere with the establishment of infection by *Leishmania*.

The role of saliva

As mentioned above, *Leishmania* promastigotes are transmitted by the bite of an infected phlebotomine. During blood feeding, the insect regurgitates and a portion of saliva is inoculated in the host dermis. In order to facilitate the blood meal, phlebotomine saliva is endowed with several substances that prevent blood clotting formation. Thus, saliva from phlebotomine sand flies (*Phlebotomus papatasi* and *Phlebotomus argentipes*) is rich in adenosine and AMP [64–66]. In addition, transcriptome analysis indicated the presence of apyrases and 5'-nucleotidases in the saliva of *Lutzomyia longipalpis* [67], *Phlebotomus perniciosus*, *P. argentipes*, *Phlebotomus ariasi* [68] and *Phlebotomus duboscqi* [69]. AMP and adenosine

inhibit platelet aggregation [70]. The presence of nucleotidases in the saliva contributes to the hydrolysis of extracellular ATP, preventing platelet aggregation and blood clot formation [71].

In addition to prevention of blood clotting, the presence of adenosine, AMP and ectonucleotidases may also influence the host immune response and facilitate the establishment of the infection. It has been demonstrated that addition of salivary gland extract together with the parasite exacerbates lesion development in the murine model of leishmaniasis [72–75]. Recent studies by Carregaro and colleagues [76] demonstrated that adenosine and AMP present in the saliva of *P. papatasi* exacerbate lesion development in *L. amazonensis* infected mice, probably due to increased IL-10 production and induction of tolerogenic dendritic cells and regulatory T cells. These effects seem to be mediated by the A_{2A} adenosine receptor. Corroborating the role of nucleotides and nucleosides of the insect saliva in the immunomodulation of the host response Katz and colleagues [64] demonstrated that *P. papatasi* saliva decreases NO production by activated macrophages. The same was not observed when the saliva of *L. longipalpis* is used. Interestingly, the levels of AMP are smaller in the saliva of *L. longipalpis* than in the saliva of *P. papatasi*. In addition, adenosine deaminase activity has been observed in the saliva of *L. longipalpis* [77] and *P. duboscqi* [78], but not in the saliva of *P. papatasi* [78]. The presence of adenosine deaminase could contribute to a decrease in adenosine concentration at the bite site thus inhibiting the modulatory effect of the saliva.

Altogether, these results indicate a strong involvement of the purinergic signaling pathway during the first moments of the inoculation of the parasite in the host dermis.

Leishmania ectonucleotidases

In addition to the effects promoted by the insect saliva, ectoenzymes present on the surface of the promastigote may also contribute to the establishment of the infection by *Leishmania*.

Leishmania and other trypanosomatids are incapable of de novo synthesis of the purine ring and thus depend on salvage pathways to synthesize purine nucleotides [79,80]. In order to obtain nucleosides from the external medium these parasites express ectonucleotidases thus enabling the external hydrolysis of nucleotides to the respective nucleosides.

Amongst the several ectonucleotidases the ectonucleoside triphosphate diphosphohydrolase (E-NTPDase – EC 3.6.1.5) and the ecto-5'-nucleotidase (EC 3.1.3.5) have been shown to exert important roles in the infection by *Leishmania*. E-NTPDase has been reported on *Leishmania tropica* [81] and *L. amazonensis* [82,83]. In addition to E-NTPDase, we also demonstrated that *L. amazonensis* presents ecto-5'-nucleotidase activity [84].

Several studies have associated the presence of E-NTPDase and ecto-5'-nucleotidase activities to the virulence of different *Leishmania* species [82,84–86]. Thus, parasites with increased ectonucleotidase activity are more capable of infecting macrophages [87] and induce larger lesions in mice [84,86]. Furthermore, we have demonstrated a positive correlation between the ATP, ADP and AMP hydrolytic activity in

Leishmania braziliensis isolates and their ability to control the establishment of the immune response in mice. More importantly, a positive correlation between the nucleotide hydrolytic activity and the severity of the clinical manifestation was also observed [85]. Similarly, *L. amazonensis* isolates with different ectonucleotidase activity show distinct capacity to survive in infected macrophages and to down-modulate nitric oxide production [87]. Interestingly, the association of ectonucleotidase activity and parasite virulence seems not to be restricted to *Leishmania*, given that it has been reported also in *Trypanosoma cruzi* [88], *Toxoplasma gondii* [89] and *Legionella pneumophila* [90]. These studies suggest that the hydrolysis of extracellular ATP to adenosine, in addition to fuel the purine salvage pathway, may also contribute to the modulation of the host immune response. In fact, we have shown that inhibition of adenosine receptors during infection by *L. braziliensis* increase the production of inflammatory cytokines and decreases tissue parasitism [84]. The fact that extracellular adenosine can modulate the immune response against the parasite suggests a potential role of ectonucleotidases in the virulence of the parasite. Corroborating this hypothesis, we and others have shown a correlation between the level of ectonucleotidase activity and expression and the virulence of the parasite, for both *Leishmania* and *T. cruzi* [86,88,91].

Interestingly, in addition to the production of extracellular adenosine, it has also been demonstrated that E-NTPDases of *T. cruzi* and *L. amazonensis* are involved in the attachment of the parasite to the host cell [83,92].

Finally, it has been shown that, although *L. amazonensis* triggers the release of neutrophil extracellular traps [10], the parasite's 3'-ectonucleotidase is capable of degrading these traps [93], enhancing even more the contribution of ectonucleotidases to the control of the innate immune response at the first moments of interaction between the parasite and the host cells.

Collectively, these results indicate that, in addition to their role in the insect saliva, ectonucleotidases from the parasite are important tools to evade the innate immune response and should be considered as potential targets for vaccine development against *Leishmania*.

Purinergic signaling and immune response

Once inoculated in the dermis of the host *Leishmania* promastigotes are phagocytized by cells present at the site of infection. Although the macrophage is the cell in which the parasite proliferates and persist to establish the infection, it has been demonstrated that *Leishmania* are also phagocytized by neutrophils and dendritic cells and this interaction can modulate the immune response to the parasite.

Neutrophils rapidly infiltrate the site of infection and phagocytize the inoculated promastigotes [8,9]. It has been proposed that interaction of the parasite with these cells induces neutrophil apoptosis. The interaction of the apoptotic neutrophil loaded [12] or not [8] with the parasite would then deactivate the macrophage, favoring parasite survival within the phagolysosome. As mentioned above, *Leishmania* also induces the release of NET (neutrophil extracellular traps) by neutrophils and may also degrade the nucleic acids present in

this structure via the action of a 3'-nucleotidase [10,93]. Although no direct proof has been yet provided, it is conceivable that nucleic acid hydrolysis may release adenosine which, by acting on the adenosine receptors A_{2A} and/or A_{2B} , could modulate the host immune response.

Dendritic cells have also been shown to be infected by *Leishmania* promastigotes [94]. This infection is capable of modulating the dendritic cell ability to present antigen to T cells thus affecting the establishment of the adaptative immune response [18,95–98]. Our group has shown that, at least part of the modulatory effect of the infection on the activation of dendritic cells (expression of CD40) is dependent on the activation of the A_{2B} adenosine receptor, mediated by the production of extracellular adenosine originated by the increased ectonucleotidase expression (CD39 and CD73) on the surface of infected cells [18]. In addition, this study also shows that inhibition of dendritic cell activation interferes with T cell proliferation which can be reverted by the blockade of the A_{2B} adenosine receptor.

Infection of macrophages by *Leishmania* is dependent on the phagocytosis of the parasite. Interestingly, an increased expression of P2X₇ receptors is observed in infected macrophages [99,100]. Due to the upregulation of the P2X₇ receptors, addition of extracellular ATP to infected macrophage cultures reduces the percentage of infected macrophages, suggesting a possible role of these receptors in the control of the parasite. The control of parasite growth by activated P2X₇ receptors seems to be mediated by the production of LTB₄ [101].

In spite of the expression of P2X₇ receptor by infected macrophages and the ability of added extracellular ATP to control parasite proliferation, *Leishmania* survives within these cells, indicating that other purinergic signaling mechanisms may exist in the infected macrophage. In fact, we have observed that macrophages infected by *L. amazonensis* express high levels of CD39 and CD73 and that inhibition of the A_{2B} adenosine receptor during infection favors the survival of the parasite (unpublished observations). Furthermore, we have demonstrated that activation of the adenosine A_{2B} receptor inhibits the production of NO and IL-12 by infected macrophages even in the presence of activating stimuli such as IFN- γ and LPS [87], allowing for the enhanced survival of the parasite. Corroborating these findings, a recent study demonstrated increased expression of the A_{2B} in monocytes from patients with visceral leishmaniasis and the possible involvement of the activation of this receptor with IL-10 production by infected cells [102].

Conclusion

Several lines of evidence indicate that purinergic signaling interferes with many aspects of the immune response against *Leishmania*. These findings provide a different approach to mechanisms of immune response evasion by the parasite similar to those being investigated in several types of tumors. Although in its initial phases, the field seems promising and may contribute to the development of new therapeutic approaches to the control of parasite progression via the activation of P2X₇ receptors or inhibition of adenosine receptors.

Conflict of interest

The authors declare that they have no competing interests.

Acknowledgements

The authors' laboratory is supported by grants from CNPq, FAPEMIG, Rede de Pesquisa em Doenças Infecciosas Humanas e Animais do Estado de Minas Gerais/FAPEMIG, UFOP. LCCA is a CNPq fellow.

REFERENCES

- [1] Herwaldt BL. Leishmaniasis. *Lancet* 1999;354:1191–9.
- [2] Kamhawi S. Phlebotomine sand flies and *Leishmania* parasites: friends or foes? *Trends Parasitol* 2006;22:439–45.
- [3] Alvar J, Velez ID, Bern C, Herrero M, Desjeux P, Cano J, et al. Leishmaniasis worldwide and global estimates of its incidence. *PLoS One* 2012;7:e35671.
- [4] Kaye P, Scott P. Leishmaniasis: complexity at the host-pathogen interface. *Nat Rev Microbiol* 2011;9:604–15.
- [5] Cecilio P, Perez-Cabezas B, Santarem N, Maciel J, Rodrigues V, Cordeiro da Silva A. Deception and manipulation: the arms of *Leishmania*, a successful parasite. *Front Immunol* 2014;5:480.
- [6] Gollob KJ, Viana AG, Dutra WO. Immunoregulation in human American leishmaniasis: balancing pathology and protection. *Parasite Immunol* 2014;36:367–76.
- [7] Gurung P, Kanneganti TD. Innate immunity against *Leishmania* infections. *Cell Microbiol* 2015;17:1286–94.
- [8] Peters NC, Egen JG, Secundino N, Debrabant A, Kimblin N, Kamhawi S, et al. In vivo imaging reveals an essential role for neutrophils in leishmaniasis transmitted by sand flies. *Science* 2008;321:970–4.
- [9] Thalhofer CJ, Chen Y, Sudan B, Love-Homan L, Wilson ME. Leukocytes infiltrate the skin and draining lymph nodes in response to the protozoan *Leishmania infantum chagasi*. *Infect Immun* 2011;79:108–17.
- [10] Guimarães-Costa AB, Nascimento MT, Froment GS, Soares RP, Morgado FN, Conceicao-Silva F, et al. *Leishmania amazonensis* promastigotes induce and are killed by neutrophil extracellular traps. *Proc Natl Acad Sci USA* 2009;106:6748–53.
- [11] Novais FO, Santiago RC, Bafica A, Khouri R, Afonso L, Borges VM, et al. Neutrophils and macrophages cooperate in host resistance against *Leishmania braziliensis* infection. *J Immunol* 2009;183:8088–98.
- [12] Laskay T, van ZG, Solbach W. Neutrophil granulocytes as host cells and transport vehicles for intracellular pathogens: apoptosis as infection-promoting factor. *Immunobiology* 2008;213:183–91.
- [13] Charmoy M, Brunner-Agten S, Aebischer D, Auderset F, Launois P, Milon G, et al. Neutrophil-derived CCL3 is essential for the rapid recruitment of dendritic cells to the site of *Leishmania major* inoculation in resistant mice. *PLoS Pathog* 2010;6:e1000755.
- [14] Berberich C, Ramirez-Pineda JR, Hambrecht C, Alber G, Skeiky YA, Moll H. Dendritic cell (DC)-based protection against an intracellular pathogen is dependent upon DC-derived IL-12 and can be induced by molecularly defined antigens. *J Immunol* 2003;170:3171–9.

- [15] Afonso LCC, Scharton TM, Vieira LQ, Wysocka M, Trinchieri G, Scott P. The adjuvant effect of Interleukin-12 in a vaccine against *Leishmania major*. *Science* 1994;263:235–7.
- [16] Heinzel FP, Schoenhaut DS, Rerko RM, Rosser LE, Gately MK. Recombinant Interleukin 12 cures mice infected with *Leishmania major*. *J Exp Med* 1993;177:1505–9.
- [17] Silveira FT, Lainson R, Corbett CE. Clinical and immunopathological spectrum of American cutaneous leishmaniasis with special reference to the disease in Amazonian Brazil: a review. *Mem Inst Oswaldo Cruz* 2004;99:239–51.
- [18] Figueiredo AB, Serafim TD, Marques-da-Silva EA, Meyer-Fernandes JR, Afonso LC. *Leishmania amazonensis* impairs DC function by inhibiting CD40 expression via A2B adenosine receptor activation. *Eur J Immunol* 2012;42:1203–15.
- [19] Olivier M, Gregory DJ, Forget G. Subversion mechanisms by which *Leishmania* parasites can escape the host immune response: a signaling point of view. *Clin Microbiol Rev* 2005;18:293–305.
- [20] Perrella Balestieri FM, Pires Queiroz AR, Scavone C, Assis C,V, Barral-Netto M, Abrahamsohn IA. *Leishmania (L.) amazonensis*-induced inhibition of nitric oxide synthesis in host macrophages. *Microb Infect* 2002;4:23–9.
- [21] Rodrigues V, Cordeiro-da-Silva A, Laforge M, Ouassii A, Akharid K, Silvestre R, et al. Impairment of T cell function in parasitic infections. *PLoS Negl Trop Dis* 2014;8.
- [22] Mosser DM, Edelson PJ. The mouse macrophage receptor for C3bi (CR3) is a major mechanism in the phagocytosis of *Leishmania* promastigotes. *J Immunol* 1985;135:2785–9.
- [23] Mosser DM, Springer TA, Diamond MS. *Leishmania* promastigotes require opsonic complement to bind to the human leukocyte integrin Mac-1 (CD11b/CD18). *J Cell Biol* 1992;116:511–20.
- [24] Peters C, Aebischer T, Stierhof YD, Fuchs M, Overath P. The role of macrophage receptors in adhesion and uptake of *Leishmania mexicana* amastigotes. *J Cell Sci* 1995;108(Pt 12):3715–24.
- [25] Wozencraft AO, Sayers G, Blackwell JM. Macrophage type 3 complement receptors mediate serum-independent binding of *Leishmania donovani*. Detection of macrophage-derived complement on the parasite surface by immunoelectron microscopy. *J Exp Med* 1986;164:1332–7.
- [26] Mosser DM, Edelson PJ. The third component of complement (C3) is responsible for the intracellular survival of *Leishmania major*. *Nature* 1987;327:329–31.
- [27] Green SJ, Crawford RM, Hockmeyer JT, Meltzer MS, Nacy CA. *Leishmania major* amastigotes initiate the L-arginine-dependent killing mechanism in IFN- γ stimulated macrophages by induction of tumor necrosis factor. *J Immunol* 1990;145:4290–7.
- [28] Mougneau E, Bihl F, Glaichenhaus N. Cell biology and immunology of *Leishmania*. *Immunol Rev* 2011;240:286–96.
- [29] Scharton-Kersten T, Scott P. The role of the innate immune response in Th1 cell development following *Leishmania major* infection. *J Leukoc Biol* 1995;57:515–22.
- [30] Scott P. IFN- γ modulates the early development of Th1 and Th2 responses in a murine model of cutaneous leishmaniasis. *J Immunol* 1991;147:3149–55.
- [31] Belkaid Y, von Stebut E, Mendez S, Lira R, Caler E, Bertholet S, et al. CD8⁺ T cells are required for primary immunity in C57BL/6 mice following low-dose, intradermal challenge with *Leishmania major*. *J Immunol* 2002;168:3992–4000.
- [32] Liese J, Schleicher U, Bogdan C. The innate immune response against *Leishmania* parasites. *Immunobiology* 2008;213:377–87.
- [33] Sacks DL. The structure and function of the surface lipophosphoglycan on different developmental stages of *Leishmania* promastigotes. *Infect Agents Dis* 1992;1:200–6.
- [34] Jones DE, Ackermann MR, Wille U, Hunter CA, Scott P. Early enhanced Th1 response after *Leishmania amazonensis* infection of C57BL/6 interleukin-10-deficient mice does not lead to resolution of infection. *Infect Immun* 2002;70:2151–8.
- [35] Di Virgilio F. Purinergic mechanism in the immune system: a signal of danger for dendritic cells. *Purinergic Signal* 2005;1:205–9.
- [36] Elliott MR, Chekeni FB, Trampont PC, Lazarowski ER, Kadl A, Walk SF, et al. Nucleotides released by apoptotic cells act as a find-me signal to promote phagocytic clearance. *Nature* 2009;461:282–6.
- [37] Idzko M, Hammad H, Van Nimwegen M, Kool M, Willart MA, Muskens F, et al. Extracellular ATP triggers and maintains asthmatic airway inflammation by activating dendritic cells. *Nat Med* 2007;13:913–9.
- [38] La Sala A, Ferrari D, Di Virgilio F, Idzko M, Norgauer J, Girolomoni G. Alerting and tuning the immune response by extracellular nucleotides. *J Leukoc Biol* 2003;73:339–43.
- [39] Praetorius HA, Leipziger J. ATP release from non-excitable cells. *Purinergic Signal* 2009;5:433–46.
- [40] Asgari E, Le Friec G, Yamamoto H, Perucha E, Sacks SS, Kohl J, et al. C3a modulates IL-1 β secretion in human monocytes by regulating ATP efflux and subsequent NLRP3 inflammasome activation. *Blood* 2013;122:3473–81.
- [41] Ayna G, Krysko DV, Kaczmarek A, Petrovski G, Vandenabeele P, Fesus L. ATP release from dying autophagic cells and their phagocytosis are crucial for inflammasome activation in macrophages. *PLoS One* 2012;7:e40069.
- [42] Junger WG. Immune cell regulation by autocrine purinergic signalling. *Nat Rev Immunol* 2011;11:201–12.
- [43] Murphy JK, Livingston FR, Gozal E, Torres M, Forman HJ. Stimulation of the rat alveolar macrophage respiratory burst by extracellular adenine nucleotides. *Am J Respir Cell Mol Biol* 1993;9:505–10.
- [44] Schnurr M, Then F, Galambos P, Scholz C, Siegmund B, Endres S, et al. Extracellular ATP and TNF- α synergize in the activation and maturation of human dendritic cells. *J Immunol* 2000;165:4704–9.
- [45] Trautmann A. Extracellular ATP in the immune system: more than just a “danger signal”. *Sci Signal* 2009;2:pe6.
- [46] Abbracchio MP, Burnstock G. Purinoceptors: are there families of P2X and P2Y purinoceptors? *Pharmacol Ther* 1994;64:445–75.
- [47] Bours MJ, Swennen EL, Di Virgilio F, Cronstein BN, Dagnelie PC. Adenosine 5'-triphosphate and adenosine as endogenous signaling molecules in immunity and inflammation. *Pharmacol Ther* 2006;112:358–404.
- [48] Burnstock G, Kennedy C. Is there a basis for distinguishing two types of P2-purinoceptor? *General Pharmacol* 1985;16:433–40.
- [49] Ferrari D, Pizzirani C, Adinolfi E, Lemoli RM, Curti A, Idzko M, et al. The P2X7 receptor: a key player in IL-1 processing and release. *J Immunol* 2006;176:3877–83.
- [50] He Y, Franchi L, Nunez G. TLR agonists stimulate Nlrp3-dependent IL-1 β production independently of the purinergic P2X7 receptor in dendritic cells and in vivo. *J Immunol* 2013;190:334–9.
- [51] Pelegrin P, Barroso-Gutierrez C, Surprenant A. P2X7 receptor differentially couples to distinct release pathways for IL-1 β in mouse macrophage. *J Immunol* 2008;180:7147–57.
- [52] Robson SC, Seigny J, Zimmermann H. The E-NTPDase family of ectonucleotidases: structure function

- relationships and pathophysiological significance. *Purinergic Signal* 2006;2:409–30.
- [53] Zimmermann H. Extracellular metabolism of ATP and other nucleotides. *Naunyn Schmiedeb Arch Pharmacol* 2000;362:299–309.
- [54] Hagberg H, Andersson P, Lacarewicz J, Jacobson I, Butcher S, Sandberg M. Extracellular adenosine, inosine, hypoxanthine, and xanthine in relation to tissue nucleotides and purines in rat striatum during transient ischemia. *J Neurochem* 1987;49:227–31.
- [55] Schulte G, Fredholm BB. Signalling from adenosine receptors to mitogen-activated protein kinases. *Cell Signal* 2003;15:813–27.
- [56] Zetterstrom T, Vernet L, Ungerstedt U, Tossman U, Jonzon B, Fredholm BB. Purine levels in the intact rat brain. Studies with an implanted perfused hollow fibre. *Neurosci Lett* 1982;29:111–5.
- [57] Fredholm BB, IJzerman AP, Jacobson KA, Klotz KN, Linden J. International Union of Pharmacology. XXV. Nomenclature and classification of adenosine receptors. *Pharmacol Rev* 2001;53:527–52.
- [58] Antonioli L, Blandizzi C, Pacher P, Hasko G. Immunity, inflammation and cancer: a leading role for adenosine. *Nat Rev Cancer* 2013;13:842–57.
- [59] Cekic C, Linden J. Purinergic regulation of the immune system. *Nature Rev Immunol* 2016;16:177–92.
- [60] Csoka B, Selmecky Z, Koscsó B, Nemeth ZH, Pacher P, Murray PJ, et al. Adenosine promotes alternative macrophage activation via A2A and A2B receptors. *FASEB J* 2012;26:376–86.
- [61] Hasko G, Kuhel DG, Chen JF, Schwarzschild MA, Deitch EA, Mabley JG, et al. Adenosine inhibits IL-12 and TNF- α production via adenosine A_{2A} receptor-dependent and independent mechanisms. *FASEB J* 2000;14:2065–74.
- [62] Panther E, Idzko M, Herouy Y, Rheinen H, Gebicke-Haerter PJ, Mrowietz U, et al. Expression and function of adenosine receptors in human dendritic cells. *FASEB J* 2001;15:1963–70.
- [63] Wilson JM, Ross WG, Agbai ON, Frazier R, Figler RA, Rieger J, et al. The A2B adenosine receptor impairs the maturation and immunogenicity of dendritic cells. *J Immunol* 2009;182:4616–23.
- [64] Katz O, Waitumbi JN, Zer R, Warburg A. Adenosine, AMP, and protein phosphatase activity in sandfly saliva. *Am J Trop Med Hyg* 2000;62:145–50.
- [65] Ribeiro JM, Katz O, Pannell LK, Waitumbi J, Warburg A. Salivary glands of the sand fly *Phlebotomus papatasi* contain pharmacologically active amounts of adenosine and 5'-AMP. *J Exp Biol* 1999;202:1551–9.
- [66] Ribeiro JM, Modi G. The salivary adenosine/AMP content of *Phlebotomus argentipes* Annandale and Brunetti, the main vector of human kala-azar. *J Parasitol* 2001;87:915–7.
- [67] Valenzuela JG, Garfield M, Rowton ED, Pham VM. Identification of the most abundant secreted proteins from the salivary glands of the sand fly *Lutzomyia longipalpis*, vector of *Leishmania chagasi*. *J Exp Biol* 2004;207:3717–29.
- [68] Anderson JM, Oliveira F, Kamhawi S, Mans BJ, Reynoso D, Seitz AE, et al. Comparative salivary gland transcriptomics of sandfly vectors of visceral leishmaniasis. *BMC Genomics* 2006;7:52.
- [69] Hamasaki R, Kato H, Terayama Y, Iwata H, Valenzuela JG. Functional characterization of a salivary apyrase from the sand fly, *Phlebotomus duboscqi*, a vector of *Leishmania major*. *J Insect Physiol* 2009;55:1044–9.
- [70] Vial C, Rolf MG, Mahaut-Smith MP, Evans RJ. A study of P2X1 receptor function in murine megakaryocytes and human platelets reveals synergy with P2Y receptors. *Br J Pharmacol* 2002;135:363–72.
- [71] Charlab R, Valenzuela JG, Rowton ED, Ribeiro JM. Toward an understanding of the biochemical and pharmacological complexity of the saliva of a hematophagous sand fly *Lutzomyia longipalpis*. *Proc Natl Acad Sci USA* 1999;96:15155–60.
- [72] Lima HC, Titus RG. Effects of sand fly vector saliva on development of cutaneous lesions and the immune response to *Leishmania braziliensis* in BALB/c mice. *Infect Immun* 1996;64:5442–5.
- [73] Mbow ML, Bleyenbergh JA, Hall LR, Titus RG. *Phlebotomus papatasi* sand fly salivary gland lysate down-regulates a Th1, but up-regulates a Th2, response in mice infected with *Leishmania major*. *J Immunol* 1998;161:5571–7.
- [74] Morris RV, Shoemaker CB, David JR, Lanzaro GC, Titus RG. Sandfly maxadilan exacerbates infection with *Leishmania major* and vaccinating against it protects against *L. major* infection. *J Immunol* 2001;167:5226–30.
- [75] Norsworthy NB, Sun J, Elnaïem D, Lanzaro G, Soong L. Sand fly saliva enhances *Leishmania amazonensis* infection by modulating interleukin-10 production. *Infect Immun* 2004;72:1240–7.
- [76] Carregaro V, Ribeiro JM, Valenzuela JG, Souza-Junior DL, Costa DL, Oliveira CJ, et al. Nucleosides present on phlebotomine saliva induce immunosuppression and promote the infection establishment. *PLoS Negl Trop Dis* 2015;9:e0003600.
- [77] Charlab R, Rowton ED, Ribeiro JM. The salivary adenosine deaminase from the sand fly *Lutzomyia longipalpis*. *Exp Parasitol* 2000;95:45–53.
- [78] Kato H, Jochim RC, Lawyer PG, Valenzuela JG. Identification and characterization of a salivary adenosine deaminase from the sand fly *Phlebotomus duboscqi*, the vector of *Leishmania major* in sub-Saharan Africa. *J Exp Biol* 2007;210:733–40.
- [79] Landfear SM, Ullman B, Carter NS, Sanchez MA. Nucleoside and nucleobase transporters in parasitic protozoa. *Eukaryot Cell* 2004;3:245–54.
- [80] Marr JJ, Berens RL, Nelson DJ. Purine metabolism in *Leishmania donovani* and *Leishmania braziliensis*. *Biochim Biophys Acta* 1978;544:360–71.
- [81] Peres-Sampaio CE, Palumbo ST, Meyer-Fernandes JR. An ecto-ATPase activity present in *Leishmania tropica* stimulated by dextran sulfate. *Z Naturforsch C* 2001;56:820–5.
- [82] Berredo-Pinho M, Peres-Sampaio CE, Chrispim PP, Belmont-Firpo R, Lemos AP, Martiny A, et al. A Mg-dependent ecto-ATPase in *Leishmania amazonensis* and its possible role in adenosine acquisition and virulence. *Arch Biochem Biophys* 2001;391:16–24.
- [83] Pinheiro CM, Martins-Duarte ES, Ferraro RB, Fonseca De Souza AL, Gomes MT, Lopes AH, et al. *Leishmania amazonensis*: biological and biochemical characterization of ecto-nucleoside triphosphate diphosphohydrolase activities. *Exp Parasitol* 2006;114:16–25.
- [84] Marques-da-Silva EdA, de Oliveira JC, Figueiredo AB, Lima Jr DdS, Carneiro CM, Fietto JLR, et al. Extracellular nucleotide metabolism in *Leishmania*: influence of adenosine in the establishment of infection. *Microb Infect* 2008;10:850–7.
- [85] Leite PM, Gomes RS, Figueiredo AB, Serafim TD, Tafuri WL, de Souza CC, et al. Ecto-nucleotidase activities of promastigotes from *Leishmania (Viannia) braziliensis* relates to parasite infectivity and disease clinical outcome. *PLoS Negl Trop Dis* 2012;6:e1850.
- [86] Souza MC, Assis EA, Gomes RS, Marques-da-Silva EA, Melo MN, Fietto JL, et al. The influence of ecto-nucleotidases

- on *Leishmania amazonensis* infection and immune response in C57B/6 mice. *Acta Trop* 2010;115:262–9.
- [87] Gomes RS, de Carvalho LC, de Souza Vasconcellos R, Fietto JL, Afonso LC. E-NTPDase (ecto-nucleoside triphosphate diphosphohydrolase) of *Leishmania amazonensis* inhibits macrophage activation. *Microbes Infect* 2015;17:295–303.
- [88] Santos RF, Possa MA, Bastos MS, Guedes PM, Almeida MR, DeMarco R, et al. Influence of ecto-nucleoside triphosphate diphosphohydrolase activity on trypanosoma cruzi infectivity and virulence. *PLoS Negl Trop Dis* 2009;3:e387.
- [89] Asai T, Miura S, Sibley LD, Okabayashi H, Takeuchi T. Biochemical and molecular characterization of nucleoside triphosphate hydrolase isozymes from the parasitic protozoan *Toxoplasma gondii*. *J Biol Chem* 1995;270:11391–7.
- [90] Sansom FM, Newton HJ, Crikis S, Cianciotto NP, Cowan PJ, d'Apice AJ, et al. A bacterial ecto-triphosphate diphosphohydrolase similar to human CD39 is essential for intracellular multiplication of *Legionella pneumophila*. *Cell Microbiol* 2007;9:1922–35.
- [91] Paletta-Silva R, Meyer-Fernandes JR. Adenosine and immune imbalance in visceral leishmaniasis: the possible role of ectonucleotidases. *J Trop Med* 2012;2012:650874.
- [92] Vasconcellos Rde S, Mariotini-Moura C, Gomes RS, Serafim TD, Firmino Rde C, Silva EBM, et al. *Leishmania infantum* ecto-nucleoside triphosphate diphosphohydrolase-2 is an apyrase involved in macrophage infection and expressed in infected dogs. *PLoS Negl Trop Dis* 2014;8:e3309.
- [93] Guimaraes-Costa AB, Desouza-Vieira TS, Paletta-Silva R, Freitas-Mesquita AL, Meyer-Fernandes JR, Saraiva EM. 3'-Nucleotidase/nuclease activity allows *Leishmania* parasites to escape killing by neutrophil extracellular traps. *Infect Immun* 2014;82:1732–40.
- [94] Brandonisio O, Spinelli R, Pepe M. Dendritic cells in *Leishmania* infection. *Microb Infect* 2004;6:1402–9.
- [95] Prina E, Abdi SZ, Lebastard M, Perret E, Winter N, Antoine JC. Dendritic cells as host cells for the promastigote and amastigote stages of *Leishmania amazonensis*: the role of opsonins in parasite uptake and dendritic cell maturation. *J Cell Sci* 2004;117:315–25.
- [96] Qi H, Popov V, Soong L. *Leishmania amazonensis*-dendritic cell interactions in vitro and the priming of parasite-specific CD4⁺ T cells in vivo. *J Immunol* 2001;167:4534–42.
- [97] Vasquez RE, Xin L, Soong L. Effects of CXCL10 on dendritic cell and CD4⁺ T-cell functions during *Leishmania amazonensis* infection. *Infect Immun* 2008;76:161–9.
- [98] Xin L, Li K, Soong L. Down-regulation of dendritic cell signaling pathways by *Leishmania amazonensis* amastigotes. *Mol Immunol* 2008;45:3371–82.
- [99] Chaves SP, Torres-Santos EC, Marques C, Figliuolo VR, Persechini PM, Coutinho-Silva R, et al. Modulation of P2X₇ purinergic receptor in macrophages by *Leishmania amazonensis* and its role in parasite elimination. *Microb Infect* 2009;11:842–9.
- [100] Coutinho-Silva R, Correa G, Sater AA, Ojcius DM. The P2X₇ receptor and intracellular pathogens: a continuing struggle. *Purinergic Signal* 2009;5:197–204.
- [101] Chaves MM, Marques-da-Silva C, Monteiro AP, Canetti C, Coutinho-Silva R. Leukotriene B₄ modulates P2X₇ receptor-mediated *Leishmania amazonensis* elimination in murine macrophages. *J Immunol* 2014;192:4765–73.
- [102] Vijayamahantesh, Amit A, Kumar S, Dikhit MR, Jha PK, Singh AK, et al. Up regulation of A2B adenosine receptor on monocytes are crucially required for immune pathogenicity in Indian patients exposed to *Leishmania donovani*. *Cytokine* 2016;79:38–44.